# Botox treatment for Migraine

Botox is a treatment for chronic migraine that can reduce how often you have migraine attacks and how severe they are.

#### What is Botox?

Botox is a type of nerve toxin that paralyses muscles. It comes from bacteria called Clostridium botulinum. It was found to be effective for people with migraine while being used in cosmetic treatment. People who had Botox experienced fewer headaches.

Doctors think it works by blocking chemicals called neurotransmitters that carry pain signals from within your brain. The same toxin is used to treat other health conditions such as cerebral palsy, bladder pain, lower back pain, neuropathic pain and stroke.

### When is Botox used?

Botox is approved for use on the NHS for the treatment of chronic migraine in adults. Chronic migraine is defined as having at least 15 headache days a month, with at least eight of those featuring migraine symptoms.

It is not recommended for episodic migraine (when attacks are less frequent) and other types of headache.

#### Who can have Botox?

Botox is only available on prescription based on certain criteria.

In England, Wales and Northern
Ireland, it is recommended for people with
chronic migraine who have already tried at least
three other preventive treatments that haven't
been effective. In Scotland, Botox is approved
for chronic migraine when people have failed to
respond to three oral preventive treatments and
any medication overuse has been managed.

If you think you are eligible for Botox and would like to be considered for treatment you can ask for a referral to a specialist. The specialist will assess your history and symptoms to make sure Botox is an appropriate treatment option for you. You will usually be asked to keep a 'headache diary' to monitor the treatment's effectiveness.

If Botox doesn't work for you, or stops working, your specialist will discuss other possible treatments with you. Our factsheets on medication and treatment options outline some of these.



#### Will it work for me?

Botox aims to reduce how often you have migraine attacks and how severe they are.

Most people have at least two treatment cycles before deciding if Botox works effectively for them. A good response to Botox is usually a 30–50% reduction in how many headaches you have. Doctors will also consider any other improvements you experience. Some people notice an improvement in their quality of life with Botox even if they don't have a big reduction in headache days.

Botox is well tolerated by most people. The most common side effects include neck pain, muscular weakness and drooping eyelids. These side effects are temporary because Botox wears off over time.

## How is Botox given?

The guidelines recommend
Botox is given as a series of
31–39 small injections. These
are given under the skin or into
the muscles in and around the
forehead, above the ears and in the neck. The
person doing your treatment will have been
trained to provide Botox for chronic migraine.

Injections are given every 12 weeks. Botox is usually given until your migraine frequency has changed to episodic migraine for three months in a row or there has been a significant improvement in your quality of life.

You may be asked to fill in regular questionnaires so improvements can be monitored. Botox may be stopped if it doesn't treat your migraine effectively enough.

The Migraine Trust can be contacted on **0808 802 0066** (Monday to Friday, 10am to 4pm) or through our contact form and Live chat service found on our website at **migrainetrust.org** or by emailing **info@migrainetrust.org**.



4th Floor, Mitre House, 44-46 Fleet Street, London EC4Y 1BN. Registered charity in England and Wales (1081300) and Scotland (SC042911). A company limited by guarantee registered in England (3996448).